

One-loop derivation of the Wilson polygon - MHV amplitude duality

A. Gorsky^{1,4} and A. Zhiboedov^{1,2,3}

¹ *Institute of Theoretical and Experimental Physics, Moscow, Russia,*

² *Joint Institute for Nuclear Research, Bogoliubov Laboratory of Theoretical Physics,
Dubna, Russia,*

³ *Moscow State University, Physics Department, Moscow, Russia,*

⁴ *FTPI, University of Minnesota.*

Abstract

We discuss the origin of the Wilson polygon - MHV amplitude duality at the perturbative level. It is shown that the duality for the MHV amplitudes at one-loop level can be proven upon the peculiar change of variables in Feynman parametrization and the use of the relation between Feynman integrals at the different space-time dimensions. Some generalization of the duality which implies the insertion of the particular vertex operator at the Wilson triangle is found for the 3-point function. We discuss analytical structure of Wilson loop diagrams and present the corresponding Landau equations. The geometrical interpretation of the loop diagram in terms of the hyperbolic geometry is discussed.

1 Introduction

The clarification of the geometrical structure behind the perturbation theory in SYM which would provide the way of summation of the series remains the challenging problem. During the last years two novel ideas concerning these issues have been developed. It was demonstrated in [1] that the important localization phenomena happens for the perturbative amplitude in the twistor space. On the other hand the stringy calculation of the amplitudes [2] suggested the hidden duality between the amplitudes in $\mathcal{N} = 4$ SYM and the Wilson polygon built from the light-like momenta of the external gluons. It is important to note that amplitudes look like to be mapped to ordinary position space Wilson loop. A connection between amplitudes and momentum space Wilson loops was investigated in [3].

This duality has been checked at one [4, 5] and two loops [6, 7] in the perturbative theory and has the chance to be all-loop exact (see [11] for the review). During this development it was also realized that the important dual superconformal symmetry is under the carpet which was clarified both in the weak coupling [8, 9] and strong coupling sides [10] of the correspondence. The dual superconformal symmetry was argued to be the consequence of the fermionic T-duality in the stringy sigma model [12] and the combination of the usual superconformal and dual superconformal symmetries implies the Yangian symmetry in the perturbative $\mathcal{N} = 4$ SYM theory [13].

In spite of the impressive progress many key issues are still to be clarified. In this paper we shall focus on the origin of the Wilson polygon- MHV amplitude duality which shall be analyzed at the one-loop level. We shall try to get the precise mapping between the one-loop diagram for MHV amplitude and the one-loop correction to the Wilson polygon. It turns out that upon the proper change of variables in the Feynman parametrization of the loop integral for the six-dimensional box diagram brings it to the form of the Wilson polygon in the four dimensions. Oppositely the four dimensional box diagram can be related with the Wilson polygon in six dimensions. The IR divergences of the amplitudes get mapped into the UV divergences of the Wilson polygon. Moreover, it is seen that the MHV amplitude obeys this special property since it can be expressed in terms of the two mass-easy box diagrams only and simple change of variables we have found does not work for the non-MHV amplitudes. Using known interplay between particular $D = 6$ and $D = 4$ integrals [14, 15] the answer can be immediately presented in terms of the finite part of the $D = 4$ two-mass easy box.

The loop amplitudes can be calculated via dispersion relations hence the duality implies that some version of the imaginary part calculations can be formulated for the loop corrections to the Wilson polygon as well. To this aim we shall slightly generalize the cut technique for the loop diagrams and shall argue that on the Wilson polygon side the dispersion calculation corresponds to the cutting of the Wilson polygon into

the several pieces and the following gluing with the insertion of particular operators. We shall also comment on the Landau equations for the singularities on the Wilson polygon side.

It is natural to search for more natural geometry behind the one-loop calculation which would shed additional light on the duality under discussion. Let us first comment on the previous studies of this issue. The one-loop correlation functions suggest the natural emergence of the AdS type geometry in three-point [17] and four-point functions [18]. Similar hyperbolic structure is also clearly seen in the one-loop effective action in the constant external field [19]. In both cases the Schwinger parametrization of the loop integral plays the crucial role. In particular for the three-point function the combination of the Schwinger parameters plays the role of the radial coordinate in the AdS_5 [17] while in the effective action case similar identification emerges in the AdS_3 submanifold [19].

The geometry behind the BDS formula [23] emerging upon summation over the loops has been suggested in [24] and the corresponding fermionic representation which supports the hidden integrability has been found. The key point is that there is natural playground for the topological strings both in the A model with the Kahler gravity and B model involving KS gravity on the moduli space of the complex structures. The both complex and Kähler types of moduli are provided by the kinematical invariants of the scattering particles.

In this paper we shall mention geometrical aspect of the one loop calculation. based on the observation of [20] related with the Kahler moduli side. It was found in [20] that the one loop box integral counts the hyperbolic volume of the 3d manifold in the space of Feynman parameters. Contrary to the Gopakumar's approach when the four-point function is treated differently from the three-point function in this approach they are considered on the equal footing. Since the 3d hyperbolic manifolds emerge naturally as the knot complements we shall make some links with the Chern-Simons calculation with the inserted Wilson loop.

The paper is organized as follows. In Section 2 we review the duality between the MHV amplitudes in $\mathcal{N} = 4$ theory and the Wilson polygon. In Section 3 we briefly explain the relevant hyperbolic geometry behind the one-loop calculations. Section 4 is devoted to the explicit derivation of the duality for MHV amplitude at the one loop level. In Section 5 we provide the simplified example of the duality for the 3-point function which involves the vertex operator on the Wilson polygon side. In Sections 6 we consider some aspects of the unitarity calculation of the Wilson polygons. Section 7 is devoted to the comments concerning the relation of the divergent contributions with the hyperbolic geometry of one-loop diagrams. In the last Section we shall collect our observations and mention the open problems.

2 The connection between Wilson polygons and MHV amplitudes

In this section we briefly review the conjectured duality between the loop amplitudes in $\mathcal{N} = 4$ theories and Wilson polygons built from the external momenta (see [11] for review).

Precisely, it was conjectured in [2] that any MHV N -leg amplitude follows from the vacuum expectation value of the Wilson loop of the special form

$$\frac{A_{all-loop}^{MHV}}{A_{tree}^{MHV}} = \langle W(p_1, p_2, \dots, p_N) \rangle \quad (1)$$

where the closed Wilson loop polygon has light-like momenta at the edges and vertices at x_i . Its closeness is provided by the total momentum conservation. At the strong coupling limit both Wilson polygon as well as the MHV amplitude are calculated in the sigma model approach.

At weak coupling to check this polygon- MHV amplitude duality one considers the expansion of the Wilson polygon in the YM coupling treating Wilson loop as placed in the coordinate space. In its weaker form the duality takes form

$$Fin[\frac{A_{all-loop}^{MHV}}{A_{tree}^{MHV}}] = Fin[\langle W(p_1, p_2, \dots, p_N) \rangle] \quad (2)$$

The perfect matching of the Wilson loop and amplitudes has been found for one [4, 5] and two loop answers up to six external legs [6, 7]. Moreover, it was demonstrated that the anomalous Ward identities for the special conformal transformations of the form

$$K^\nu W(x_1, \dots, x_N) = \sum_{i=1}^n (2x_i^\nu x_i \partial_i - x_i^2 \partial_i^\nu) W(x_1, \dots, x_N) = \frac{1}{2} \Gamma_{cusp} \sum_{i=1}^n \ln \frac{x_{i,i+2}^2}{x_{i-1,i+1}^2} x_{i,i+1}^\nu \quad (3)$$

where Γ_{cusp} is the cusp anomalous dimension, fix the answer up to four external legs

The anomalous Ward identities can be applied both to the amplitudes and the Wilson polygons, however, starting with six external legs the Ward identity allows the arbitrary function of the conformal ratios, which can not be fixed by the superconformal group arguments.

There is some specifics concerning the loop MHV amplitudes. The one-loop Wilson loop diagram with the arbitrary number of external legs can be mapped to the finite part of two-mass easy box which is the main building block of the answer. The generalization of the duality to the non-MHV amplitudes turns out to be nontrivial

issue. In particular it is known [37] that the NMHV loop amplitude involves 3 mass box diagrams as well and harder diagrams are relevant for the $N^k MHV$ amplitudes. No recipe for the duality beyond the MHV case has been formulated yet.

It was demonstrated that the unitarity approach is fruitful for the description of the loop amplitudes. General planar color-ordered one-loop scattering super-amplitude can be written in the following way

$$\mathcal{A}_{n;1} = i(2\pi)^4 \delta^4(p) \sum (\mathcal{C}^{4m} I^{4m} + \mathcal{C}^{3m} I^{3m} + \mathcal{C}^{2mh} I^{2mh} + \mathcal{C}^{2me} I^{2me} + \mathcal{C}^{1m} I^{1m}) \quad (4)$$

where I 's are the scalar-box integrals with the corresponding number of legs being off-shell.

The only thing one needs to calculate for given amplitude is the coefficients, which can be done in terms of quadruple cuts. The general form of the $\mathcal{C}^m$ takes the form

$$\mathcal{C}^m = \delta^8 \left(\sum_{i=1}^n \lambda_i \eta_i \right) [\mathcal{P}_{n;1}^{(0),m} + \mathcal{P}_{n;1}^{(4),m} + \dots + \mathcal{P}_{n;1}^{(4n-16),m}] \quad (5)$$

where $\mathcal{P}_{n;1}^{(4k),m}$'s are homogenous polynomials of degree $4k$ in Grassmann variables.

One-loop MHV super-amplitude takes the following form

$$\mathcal{A}_{n;1}^{MHV} = i(2\pi)^4 \delta^4(p) \frac{\delta^8(\sum_{i=1}^n \lambda_i \eta_i)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle} \left[\sum_{s=3}^{n-1} I_{1,2,s,s+1}^{2me} \Delta_{1,2,s,s+1} + cyclic \right] \quad (6)$$

where $\Delta_{r,t,s,s+1} = -\frac{1}{2}[x_{sr}^2 x_{s+1t}^2 - x_{s+1r}^2 x_{st}^2]$. The answer is fully defined by two-mass easy boxes.

The general one-loop NMHV amplitude has more complicated structure, namely

$$\mathcal{A}_{n;1}^{NMHV} = \mathcal{A}_{n;0}^{MHV} \left[\sum_{p,q,r=1}^n R_{pqr} \left(1 + \frac{\lambda}{8\pi^2} V_{pqr} + O(\varepsilon) \right) \right] \quad (7)$$

where two-mass hard and three-mass boxes are involved, R_{pqr} are dual superconformal and V_{pqr} are dual conformal invariants.

3 Hyperbolic geometry of one loop

In what follows it will be useful to utilize the geometrical picture behind the one-loop calculations which we shall review following [20]. Let us explain first the explicit map of the box diagram to the hyperbolic volume of the particular simplex build from the kinematical invariants of the external momenta. To this aim introduce the Feynman

parametrization of the internal generically massive propagators with the parameters α_i . If one considers the one-loop N-point function with the external momenta p_i in D space-time dimensions it can be brought into the usual form

$$J(D, p_1, \dots, p_N) \propto \int_0^1 \dots \int_0^1 \prod d\alpha_i \delta(\sum \alpha_i - 1) [\sum \alpha_i^2 m_i^2 + \sum_{j < l} \alpha_i \alpha_j m_j m_l C_{il}]^{D/2-N} \quad (8)$$

where

$$C_{jl} = \frac{m_i^2 + m_l^2 - k_{jl}^2}{2m_j m_i}, \quad k_{ij} = p_i - p_j \quad (9)$$

and m_i is the mass in the i-th propagator.

It is possible [20] to organize for the generic one-loop diagram the N dimensional simplex defined as follows. First introduce the N mass vectors $m_i a_i$, where a_i are the unit vectors. The length of the side connecting the i-th and j-th mass vectors is $\sqrt{k_{ij}}$ that is one can define the momentum side of the simplex. Therefore the N -dimensional simplex involves $\frac{N(N+1)}{2}$ sides including N mass sides as well as $\frac{N(N-1)}{2}$ momentum sides. At each vertex N sides meet and at all vertices but one there are one mass side and $(N-1)$ momentum sides. The volume of such N -dimensional simplex is given as follows

$$V^{(N)} = \frac{(\prod m_i) \sqrt{\det C}}{N!} \quad (10)$$

There are $(N+1)$ hypersurfaces of dimension $(N-1)$ one of which contains only momentum sides and can be related with the massless N-point function.

It is convenient to make the change of variables which transforms the loop integral into the following form

$$J(D, p_1, \dots, p_N) \propto \prod m_i^{-1} \int_0^\infty \dots \int_0^\infty \prod d\alpha_i \delta(\alpha^T C \alpha - 1) (\sum \frac{\alpha_i}{m_i})^{N-D} \quad (11)$$

that is integration now is over the quadrics in the space of the Feynman parameters. It is useful to introduce the content of the N -dimensional solid angle $\Omega^{(N)}$ subtended by the hypersurfaces at the mass meeting point. It turns out that $\Omega^{(N)}$ coincides with the content of the $(N-1)$ dimensional simplex in the hyperbolic space whose sides are equal to the hyperbolic angles τ_{ij} defined at small masses as follows

$$C_{ij} = \cosh \tau_{ij} \quad (12)$$

Then the integral for the case $D = N$ acquires the following form

$$J(N, p_1, \dots, p_N) = i^{1-2N} \frac{\pi^{N/2} \Gamma(N/2) \Omega^{(N)}}{N! V^{(N)}} \quad (13)$$

hence the calculation of the Feynman integral is nothing but the calculation of the hyperbolic volume in the proper space. The case $N \neq D$ can be treated similarly with some modification [20].

Let us turn now to the case of interest that is N-leg MHV amplitudes in four dimensions. The crucial point is that one-loop MHV amplitudes can be presented as the sum of the two mass-easy box diagrams. This diagrams are IR divergent that is it is useful to start with the box diagram with all off-shell particles. We have the situation with $D = N$ simplices in the hyperbolic space.

$$J(4, p_1, p_2, p_3, p_4) = \frac{2i\pi^2\Omega^{(4)}}{m_1m_2m_3m_4\sqrt{\det C}} \quad (14)$$

and since all internal propagators are massless in our case we get the ideal hyperbolic tetrahedron whose all vertices are at infinity. In the massless limit we get

$$(m_i^2m_2^2m_3^2m_4^2\det C)_{m_i \rightarrow 0} = \frac{1}{16}\lambda(k_{12}^2k_{34}^2, k_{13}^2k_{24}^2, k_{14}^2k_{23}^2) \quad (15)$$

where the Källén function $\lambda(x, y, z)$ is defined as

$$\lambda(x, y, z) = x^2 + y^2 + z^2 - 2xy - 2yz - 2zx \quad (16)$$

and $\sqrt{-\lambda}$ is just the area of the triangle with sides $\sqrt{k_{12}^2k_{34}^2}$, $\sqrt{k_{23}^2k_{41}^2}$, $\sqrt{k_{31}^2k_{24}^2}$. The hyperbolic volume of the ideal tetrahedron under consideration reads as

$$2i\Omega^{(4)} = Cl_2(\psi_{12}) + Cl_2(\psi_{13}) + Cl_2(\psi_{23}) \quad (17)$$

where the dihedral angles are defined via the kinematical invariants

$$-\cos\psi_{12} = \frac{k_{13}^2k_{24}^2 + k_{14}^2k_{23}^2 - k_{12}^2k_{34}^2}{\sqrt{k_{13}^2k_{23}^2k_{14}^2k_{43}^2}} \quad (18)$$

$$-\cos\psi_{13} = \frac{k_{14}^2k_{23}^2 + k_{12}^2k_{43}^2 - k_{13}^2k_{24}^2}{\sqrt{k_{14}^2k_{23}^2k_{12}^2k_{43}^2}} \quad (19)$$

$$-\cos\psi_{14} = \frac{k_{12}^2k_{34}^2 + k_{13}^2k_{24}^2 - k_{14}^2k_{32}^2}{\sqrt{k_{13}^2k_{24}^2k_{12}^2k_{43}^2}} \quad (20)$$

and $\psi_{12} = \psi_{34}$, $\psi_{13} = \psi_{24}$, $\psi_{14} = \psi_{32}$. The functions involved are defined as

$$Cl_2(x) = \text{Im}[Li_2(e^{ix})] = -\int_0^x dy \ln|2\sin y/2| \quad (21)$$

In the case of the two mass-easy box diagram defining the one-loop MHV amplitude the additional simplification of the kinematical invariants happens since two external particles are on the mass shell. In this case the arguments of the Li_2 function degenerates to the conformal ratios of four points. The geometrical picture behind the divergent part of the diagram will be discussed later.


Note that massless four-point box answer coincides with the three-point result which is known for a while [22]. However the geometrical object responsible for the three-point function is just the triangle. The answer for the generic three-point function is expressed in terms of the angles of the basic triangle only [20].

The appearance of the hyperbolic volume implies that the topological string approach or CS with $SL(2, C)$ group are relevant [40]. Indeed we can consider the ideal tetrahedron as the knot complement and shall calculate it via the Chern-Simons theory action with the complex group. It turns out that the choice of the particular values of the kinematical invariants corresponds to the choice of particular knot [27].

4 Derivation of the Wilson polygon - MHV amplitude duality at one loop

In this Section we shall derive the duality at one-loop level via the two step procedure. First we describe the change of variables in the space of Feynman parameters which brings the two mass-easy box diagrams into the form of the Wilson polygon in the different dimension. Then we take use of the relation between the Feynman diagrams in $D = 6$ and $D = 4$.

Let us start with the definition of general box in $D_{IR} = d_{IR} - 2\epsilon_{IR}$ dimensions.



and use notations from [16]

$$I(p_i, D_{IR}, \mu_{IR}) = -i\pi^{-\frac{D_{IR}}{2}} (\mu_{IR}^2)^{\epsilon_{IR}} \int d^{D_{IR}} l \frac{1}{l^2(l-p_1)^2(l-p_1-p_2)^2(l+p_4)^2} \quad (22)$$

$$p_i^2 = m_i^2$$


One can introduce Feynman parameters and take the integral over l which amounts to

$$I(p_i, D_{IR}, \mu_{IR}) = (\mu_{IR}^2)^{\epsilon_{IR}} \Gamma(4 - \frac{D_{IR}}{2}) \int \prod dx_i \frac{\delta(1 - x_1 - x_2 - x_3 - x_4)}{(-\Delta)^{4 - \frac{D_{IR}}{2}}} \quad (23)$$

$$\Delta = sx_1x_3 + ux_2x_4 + m_1^2x_1x_2 + m_2^2x_2x_3 + m_3^2x_3x_4 + m_4^2x_4x_1$$

where $s = (p_1 + p_2)^2$ and $u = (p_2 + p_3)^2$.

Let us focus on the two-mass easy box diagram



when $m_1^2 = m_3^2 = 0$ and therefore

$$\Delta_{2me} = sx_1x_3 + ux_2x_4 + m_2^2x_2x_3 + m_4^2x_4x_1 \quad (24)$$

Upon the following change of variables

$$\begin{aligned} x_1 &= \sigma_1(1 - \tau_1) \\ x_2 &= \sigma_1\tau_1 \\ x_3 &= \sigma_2\tau_2 \\ x_4 &= \sigma_2(1 - \tau_2) \\ \left| \frac{\partial(x_i)}{\partial(\sigma_i, \tau_i)} \right| &= \sigma_1\sigma_2 \end{aligned} \quad (25)$$


the integration over σ_i factorizes and one gets

$$\begin{aligned} I^{2me}(p_i, D_{IR}, \mu_{IR}) &= (\mu_{IR}^2)^{\epsilon_{IR}} \Gamma(4 - \frac{D_{IR}}{2}) \int d\sigma_1 d\sigma_2 \sigma_1^{\frac{D_{IR}}{2}-3} \sigma_2^{\frac{D_{IR}}{2}-3} \delta(1 - \sigma_1 - \sigma_2) \\ &\int_0^1 d\tau_1 d\tau_2 \frac{1}{(-(s + u - m_2^2 - m_4^2)\tau_1\tau_2 + (u - m_2^2)\tau_1 + (s - m_2^2)\tau_2 + m_2^2)^{4-\frac{D_{IR}}{2}}} \end{aligned} \quad (26)$$

In this expression one can observe much similarity with the Wilson loop diagram. Indeed, we will show further, that proper identification of parameters allows us to connect it with Wilson loop diagram explicitly.

It is important that the special combinations of Feynman parameters play the role of parametrization of the point in the Wilson polygon which emerges in one-loop calculation

$$W(\mathcal{C}_n) = \frac{1}{N} \text{Tr} \mathcal{P} \exp[ig \oint d\tau \dot{x}^\mu(\tau) A_\mu(x(\tau))] \quad (27)$$



We assume that $D_{UV} = d_{UV} - 2\epsilon_{UV}$, p_1 and p_3 are light-like, $x_1 = p_1\tau_1$, $x_2 = p_1 + p_2 + p_3\tau_2$ and the standard propagator in the Feynman gauge

$$G_{\mu\nu}^F(x-y) = -\eta_{\mu\nu} \frac{(\pi\mu_{UV}^2)^{\epsilon_{UV}}}{4\pi^2} \frac{\Gamma(\frac{D_{UV}}{2} - 1)}{(-(x-y)^2 + i\epsilon)^{\frac{D_{UV}}{2}-1}} \quad (28)$$

Ignoring trivial factor $\frac{g^2 C_F}{16\pi^2}$ we get the following expression for the diagram

$$I_{ij}^W(p_i, D_{UV}, \mu_{UV}) = \Gamma(\frac{D_{UV}}{2} - 1) (\pi\mu_{UV}^2)^{\epsilon_{UV}} \quad (29)$$

$$\int_0^1 d\tau_i d\tau_j \frac{m_2^2 + m_4^2 - s - u}{(-(s+u-m_2^2-m_4^2)\tau_1\tau_2 + \tau_1(s-m_2^2) + \tau_2(u-m_2^2) + m_2^2)^{\frac{D_{UV}}{2}-1}}$$

and from (26) and (29) we can make identification of the parameters to match two expressions. Namely substituting $\frac{D_{UV}}{2} - 1 = 4 - \frac{D_{IR}}{2}$ we get

$$\begin{aligned} d_{UV} + d_{IR} &= 10 \\ \epsilon_{IR} &= -\epsilon_{UV} \\ (\mu_{UV}^2\pi)^{\epsilon_{UV}} &= (\mu_{IR}^2)^{\epsilon_{IR}} \end{aligned} \quad (30)$$

and the exact correspondence reads as follows

$$\begin{aligned} I^{2me}(p_i, D_{IR}, \mu_{IR}) &= \frac{1}{m_2^2 + m_4^2 - s - u} \int_0^1 d\sigma \sigma^{2-\frac{D_{UV}}{2}} (1-\sigma)^{2-\frac{D_{UV}}{2}} I_{ij}^W(p_i, D_{UV}, \mu_{UV}) \\ &= \frac{1}{m_2^2 + m_4^2 - s - u} \frac{\Gamma(3 - \frac{D_{UV}}{2})^2}{\Gamma(6 - D_{UV})} I_{ij}^W(p_i, D_{UV}, \mu_{UV}) \end{aligned}$$

Note that it is possible to represent the expression for the Wilson polygon in the form which involves a kind of the integral over the reparametrization of the boundary

contour.

$$I^{2me}(p_i, D_{IR}, \mu_{IR}) = \frac{1}{m_2^2 + m_4^2 - s - u} \int_0^1 d\sigma I^W(\mathcal{C}(\sigma), p_i, D_{UV}, \mu_{UV})$$

$$\mathcal{C}(\sigma) : p_i \rightarrow \sqrt{\sigma(1-\sigma)} p_i$$

Such form of the answer was suggested at the strong coupling [32] when the integration over the reparametrizations of the boundary of the Wilson loop is necessary to restore the conformal invariance of the answer.

Suppose we are interested in $D_{UV} = 4 - 2\epsilon$ Wilson loop diagram. Than, using the known connection between Wilson diagram and finite part of the box, we get

$$I_{ij}^W(p_i, 4 - 2\epsilon) = (m_2^2 + m_4^2 - s - u) \frac{\Gamma(2 + 2\epsilon)}{\Gamma^2(1 + \epsilon)} I^{2me}(p_i, 6 + 2\epsilon) \quad (31)$$

$$= Fin\left[\frac{\Gamma(1 + 2\epsilon)}{\Gamma^2(1 + \epsilon)} I^{2me}(p_i, 4 + 2\epsilon) \frac{1}{2}(m_2^2 m_4^2 - su)\right]$$

and therefore the following relation provides the desired duality

$$I^{2me}(p_i, 6 + 2\epsilon) = Fin\left[\frac{I^{2me}(p_i, 4 + 2\epsilon)}{1 + 2\epsilon} \frac{(su - m_2^2 m_4^2)}{2(s + u - m_2^2 - m_4^2)}\right] \quad (32)$$

Such connection between D and $D - 2$ dimensional scalar loop integrals indeed exists and goes to the papers [38]. Here we are interested in the case of $D = 6$ two-mass easy boxes and their connection with $D = 4$ ones [15]. The formula reads as follows (see appendix A)

$$I^{2me}(6 + 2\epsilon) = \frac{1}{(1 + 2\epsilon)z_0} (I^{2me}(4 + 2\epsilon) - \sum_{i=1}^4 z_i I^{2me}(4 + 2\epsilon; 1 - \delta_{ki}))$$

where

$$z_0 = \sum_{i=1}^4 z_i = 2 \frac{s + u - m_2^2 - m_4^2}{su - m_2^2 m_4^2}$$

$$z_1 = \frac{u - m_2^2}{su - m_2^2 m_4^2}$$

$$z_2 = \frac{s - m_4^2}{su - m_2^2 m_4^2}$$

$$z_3 = \frac{u - m_4^2}{su - m_2^2 m_4^2}$$

$$z_4 = \frac{s - m_2^2}{su - m_2^2 m_4^2}$$

As can be easily seen the $\sum_{i=1}^4 z_i I^4(4+2\epsilon; 1-\delta_{ki})$ does precisely the job of taking the finite part.

As we know from the calculation of the one-loop NMHV amplitudes [28] the new ingredients emerge namely two-mass hard and three-mass boxes. Thus if one wants to extend the duality between Wilson loop and amplitudes to NMHV case one should be able to get these ingredients from Wilson loop language.

In the case of two-mass easy box the structure of function in the space of Feynman parameters space allowed us using change of variables to get the Wilson loop diagram multiplied by the simple numerical integral. We can interpret this integral as an integral over the reparameterizations of the contours. One can try to use the same approach of splitting Feynman parameters in two pairs: one pair parameterizes the contour while the second yields the standard parametrization of points where gluon propagator is attached.

For more complicated cases than two-mass easy box the factorization fails and therefore the simple geometrical interpretation does not work. Namely if we make all legs massive in the Feynman box and consider the corresponding Wilson contour the integrands in the amplitude and the Wilson loop looks as follows:

$$\begin{aligned} \Delta_W = & -(s+u-m_2^2-m_4^2)\tau_1\tau_2 + (u-m_2^2)\tau_1 + (s-m_2^2)\tau_2 + m_2^2 \\ & -m_1^2\tau_1(1-\tau_1) - m_3^2\tau_2(1-\tau_2) \end{aligned} \quad (33)$$

$$\begin{aligned} \Delta_A = & \sigma_1\sigma_2[-(s+u-m_2^2-m_4^2)\tau_1\tau_2 + (u-m_2^2)\tau_1 + (s-m_2^2)\tau_2 + m_2^2] \\ & + m_1^2\sigma_1^2\tau_1(1-\tau_1) + m_3^2\sigma_2^2\tau_2(1-\tau_2) \\ = & \sigma_1\sigma_2\Delta_W + m_1^2\sigma_1\tau_1(1-\tau_1) + m_3^2\sigma_2\tau_2(1-\tau_2) \end{aligned} \quad (34)$$

We have not found simple geometrical interpretation of transformation from Δ_W to Δ_A in terms of the reparameterizations of the Wilson contour and we can not naturally connect two-mass hard and harder boxes diagrams with Wilson diagrams for correspondent contours. That is if the connection between NMHV amplitudes and Wilson polygon-like objects exists which is expected according to the T-dual picture of $AdS_5 \times S_5$ superstring [12] than it seems to be more complicated.

5 3-point function - Wilson triangle duality


In this section we consider the example of the similar duality for the 3-point function and it will be clear how the generalization of the duality for the "two-mass

hard” diagram involves the particular vertex operator. To start with let us mention also interesting relation between the one-loop 3-point amplitude and the two-loop vacuum energy in the scalar theory. Namely if one considers the 3-point function $I(p_1^2, p_2^2, p_3^2)$ with the external virtualities p_1^2, p_2^2, p_3^2 and the two-loop vacuum energy $J(m_1^2, m_2^2, m_3^2)$ with the masses of the three internal propagators m_1^2, m_2^2, m_3^2 then the following relation holds [21]

$$I(D = 4 - 2\epsilon, p_1^2, p_2^2, p_3^2) = J(4 + 2\epsilon, m_1^2, m_2^2, m_3^2) \quad (35)$$

That is the duality discussed below can be applied both for the one-loop amplitude and the two-loop vacuum energy.

Consider the most general triangle in the massless scalar theory



In the Feynman parametrization it is equal to

$$\begin{aligned} p_1 + p_2 + p_3 &= 0 \\ I_\Delta(p_i, D_{IR}, \mu_{IR}) &= -(\mu_{IR}^2)^{\epsilon_{IR}} \Gamma(3 - \frac{D_{IR}}{2}) \int \prod dx_i \frac{\delta(1 - x_1 - x_2 - x_3)}{(-\Delta)^{3 - \frac{D_{IR}}{2}}} \\ \Delta &= m_3^2 x_1 x_2 + m_2^2 x_1 x_3 + m_1^2 x_2 x_3 \end{aligned} \quad (36)$$


and assuming $p_3^2 = 0$ we have

$$\Delta = m_2^2 x_1 x_3 + m_1^2 x_2 x_3$$

Let us make the following change of variables


$$\begin{aligned} x_1 &= \sigma(1 - \tau) \\ x_2 &= \sigma\tau \end{aligned} \quad (37)$$

which amounts to



$$I_{\Delta}(p_i, D_{IR}, \mu_{IR}) = (\mu_{IR}^2)^{\epsilon_{IR}} \Gamma(3 - \frac{D_{IR}}{2}) \int d\sigma dx_3 \sigma \frac{\delta(1 - \sigma - x_3)}{(\sigma x_3)^{3 - \frac{D_{IR}}{2}}} \int_0^1 d\tau \frac{1}{(m_2^2(1 - \tau) + m_1^2\tau)^{3 - \frac{D_{IR}}{2}}} \quad (38)$$

In Wilson-dual language we can interpret it in the following way



$$m_2^2(1 - \tau) + m_1^2\tau = (p_2 + p_3\tau)^2 \quad (39)$$

and the identification of parameters reads as follows

$$\begin{aligned} d_{UV} + d_{IR} &= 8 \\ \epsilon_{IR} &= -\epsilon_{UV} \\ (\mu_{UV}^2\pi)^{\epsilon_{UV}} &= (\mu_{IR}^2)^{\epsilon_{IR}} \end{aligned} \quad (40)$$

Therefore this diagram can be understood assuming the presence of the vertex operator

$$< Tr \mathcal{P} q^\mu A_\mu(x_b) \exp[ig \oint_{\mathcal{C}} d\tau \dot{x}^\mu(\tau) A_\mu(x(\tau))] > \quad (41)$$

where q^μ can be chosen as be arbitrary vector which is not orthogonal to p_3 in Minkowski sense, $(p_3 q) \neq 0$. This q^μ can be naturally identified with the polarization vector of correspondent external gluon. Hence we have an example of possible extension of Wilson dual side, when it becomes sensitive to polarizations of external gluons. This example provides some intuition for the possible generalization of the duality to less symmetric theories or NMHV amplitudes. Nevertheless, the problem of the interpretation the three-mass triangle and all boxes harder than two-mass easy one in terms of Wilson loop diagrams is still open.

6 Analytical structure of light-like Wilson loop

6.1 General comments

In this section we discuss the analytical structure of light-like Wilson loop. If the correspondence between MHV amplitudes and Wilson loops is true at any order of perturbation theory obviously their analytical structure namely the location of singularities, branches and discontinuities in the space of kinematic moduli should match each other. Thus there emerges two interesting problems on its own: analytical structure of perturbative light-like Wilson loop and the similar question concerning the areas in AdS_5 bounded by the light-like contour.

Here we start analysis of analytical structure of perturbative Wilson loop. Firstly, we can do it using its connection with scattering amplitudes. The fact of unitarity of QFT leads to optical theorem and allows one to take different branch cuts and develop generalized unitarity method to simplify loop computations. Using the correspondence between Wilson loops and amplitudes we can reformulate optical theorem at one loop in terms of Wilson loops.

Secondly, one can analyze the analytical structure of every Wilson diagram on its own. The systematic method of clarifying the structure of singularities of Feynman amplitudes was developed long time ago in the theory of analytic S-matrix. It can be obviously applied to Wilson loop diagrams. At one-loop level using the results of previous section we can apply Cutkosky rules to $10 - D_W$ boxes which are dual to Wilson diagrams to get the result for given diagram while at higher orders the additional arguments are required.

6.2 Landau singularities for the Wilson loop

Consideration here is parallel to [35], where excellent introduction to the problem can be found. Suppose we deal with scalar massless theory Feynman integrals in D dimensions than we have for any diagram [36]:

$$I \simeq \int_0^1 \prod d\alpha_i \delta(1 - \sum_i \alpha_i) \frac{\mathcal{U}^{N-(L+1)\frac{D}{2}}}{(-\mathcal{V})^{N-L\frac{D}{2}}} \quad (42)$$

Here: L - number of loops; N - number of propagators; α_i corresponds to the i -th propagator in the diagram of the form $\frac{1}{q_i^2}$.

$\mathcal{U} = \sum_{T \in T_1} \prod_{i \in \bar{T}} \alpha_i$ - sum over so-called 1-trees, degree L in α .

$\mathcal{V} = \sum_{T \in T_2} \prod_{i \in \bar{T}} \alpha_i (Q_T)^2$ - sum over so-called 2-trees, degree $(L+1)$ in α .

The I can be considered as the function of complex kinematical parameters and the natural question arises: where its singularities in the space of parameters are located? The answer to this question is given by the Landau equations which can be written in the following form:

$$\begin{cases} \frac{\partial \mathcal{V}}{\partial \alpha_i} = 0 & \wedge & \alpha_i = 0 \\ \mathcal{V} = 0 \end{cases}$$

The same analysis can be applied for any particular Wilson loop. If one considers the family of more simple diagrams where every propagator has one leg lying on the boundary, than consideration is in full analogy with the case of amplitudes. Namely the diagram has the following structure

$$W \simeq \int_0^1 \prod_{i=1}^{2N} d\tau_i \Theta_{Path}(x(\tau_{\sigma_1}) > x(\tau_{\sigma_2}) > \dots > x(\tau_{\sigma_{2N}})) \prod_{k=1}^{V_3} \hat{L}_k \frac{d^D z_1 d^D z_2 \dots d^D z_V}{\prod_{k=1}^N (-x_k^2)^{\frac{D}{2}-1}} \quad (43)$$

Here $\hat{L}_k$ is the differential operator independent of z_i 's, which comes from three-gluon vertexes [39]; V_3 - number of three-gluon vertexes

$$\begin{aligned} & A^{\mu_1} A^{\mu_2} A^{\mu_3} \int d^D z_k Tr[\partial_\mu (A_\nu [A^\mu, A^\nu])(z_k) \\ & \sim [\eta^{\mu_1 \mu_2} (\partial_1^{\mu_3} - \partial_2^{\mu_3}) + \eta^{\mu_2 \mu_3} (\partial_1^{\mu_1} - \partial_2^{\mu_1}) + \eta^{\mu_1 \mu_3} (\partial_1^{\mu_2} - \partial_2^{\mu_2})] G(x_1, x_2, x_3) \\ & \quad \hat{L}_k^{\mu_1 \mu_2 \mu_3} G(x_1, x_2, x_3) \end{aligned} \quad (44)$$

Than if the points (x_1, x_2, x_3) lie on the edges (y_1, y_2, y_3)

$$\hat{L}_k = \dot{y}_{1\mu_1} \dot{y}_{2\mu_2} \dot{y}_{3\mu_3} \hat{L}_k^{\mu_1 \mu_2 \mu_3} \quad (45)$$

For any ordering, there exists the change of variables of integration with the Jacobian J independent on the kinematical variables that makes the simple integration interval:

$$\int_0^1 \prod_{i=1}^{2N} d\tau_i \Theta_{Path}(\tau) \rightarrow \int_0^1 \prod_{i=1}^{2N} d\tilde{\tau}_i J(\tilde{\tau}) \quad (46)$$

If we have the following ordering along one of the edges $\int_0^1 d\tau_n \int_0^{\tau_n} d\tau_{n-1} \dots \int_0^{\tau_2} d\tau_1$.
 Than one can choose

$$\begin{aligned}
 \tau_n &= \tilde{\tau}_n \\
 \tau_{n-1} &= \tau_n \tilde{\tau}_{n-1} = \tilde{\tau}_n \tilde{\tau}_{n-1} \\
 &\dots \\
 \tau_1 &= \tau_2 \tilde{\tau}_1 = \tilde{\tau}_n \tilde{\tau}_{n-1} \dots \tilde{\tau}_1 \\
 J(\tilde{\tau}) &= \prod_{j=1}^n \tilde{\tau}_j^{-1}
 \end{aligned} \tag{47}$$

If the vertexes are absent than the Landau equations take the form

$$\begin{cases} \frac{\partial(\sum \alpha_k x_k^2)}{\partial \alpha_i} = 0 & \wedge & \alpha_i = 0 \\ \frac{\partial(\sum \alpha_k x_k^2)}{\partial \tilde{\tau}_i} = 0 & \wedge & \tilde{\tau}_i = 0 \wedge \tilde{\tau}_i = 1 \\ \sum \alpha_k x_k^2 = 0 \end{cases}$$

In the presence of vertexes we can introduce Feynman parameters

$$W \simeq \int_0^1 \prod_{i=1}^{2N} d\tilde{\tau}_i J(\tilde{\tau}) \prod_{k=1}^{V_3} \hat{L}_k \int_0^1 \prod_{k=1}^N d\alpha_k \alpha_k^{\frac{D}{2}-2} \delta(1 - \sum_i \alpha_i) \frac{d^D z_1 d^D z_2 \dots d^D z_V}{[-\sum \alpha_k x_k^2]^{N(\frac{D}{2}-1)}} \tag{48}$$

and integration over the vertex position could be done yielding the answer

$$W \simeq \int_0^1 \prod_{i=1}^{2N} d\tilde{\tau}_i J(\tilde{\tau}) \prod_{k=1}^{V_3} \hat{L}_k \int_0^1 \prod d\alpha_k \alpha_k^{\frac{D}{2}-2} \delta(1 - \sum_i \alpha_i) \frac{\mathcal{U}_W(\alpha_i)}{(-\mathcal{V}_W)^{(N-V)\frac{D}{2}-N}} \tag{49}$$

Here: V - number of vertexes; N - number of propagators; α_i corresponds to the i -th propagator in the diagram of the form $\frac{1}{(-x_i^2)^{\frac{D}{2}-1}}$; $\mathcal{U}_W$ and $\mathcal{V}_W$ - the result of the integration over the loop momenta.

Finally we get the following Landau equations

$$\begin{cases} \frac{\partial \mathcal{V}_W}{\partial \alpha_i} = 0 & \wedge & \alpha_i = 0 \\ \frac{\partial \mathcal{V}_W}{\partial \tilde{\tau}_i} = 0 & \wedge & \tilde{\tau}_i = 0 \wedge \tilde{\tau}_i = 1 \\ \mathcal{V}_W = 0 \end{cases}$$

6.3 Imaginary part of the Wilson loop at one loop

In unitary theory one can exploit the unitarity of S -matrix to get the following identity

$$S^+ S = 1$$

$$S = 1 + iT$$

$$\begin{aligned} 2Im(\mathcal{A}(in \rightarrow out)) &= -i(\mathcal{A}(in \rightarrow out) + \mathcal{A}^*(out \rightarrow in)) \\ &= \sum_{states} \mathcal{A}^*(out \rightarrow all)\mathcal{A}(in \rightarrow all) \end{aligned} \quad (50)$$

We are interested in amplitudes with n outgoing particles. In that case the RHS sum becomes the sum over state with integration over a Lorentz invariant phase space corresponding to the final particles. Pushing this statement to diagrammatic level one ends with Cutkosky rules and prescription of cutting propagators:

$$\frac{1}{k^2 + i\epsilon} \rightarrow \Theta(k_0)\delta(k^2) \quad (51)$$

It is well-known that to get the imaginary part of the diagram to given order one should sum over all possible cuts of all diagrams and over all possible intermediate states. Than one should do the integration over LIPS. The Wilson loop knows about $\mathcal{N} = 4$ SYM particle content only through correction to the gluon propagator and the vertices. At one loop level it is obviously insensitive to particle content. On the amplitudes side the cut is on the contrary essentially dependent on the particle content and tree-level amplitude even at one loop. According to the strong version of the correspondence which is true at one loop


$$\mathcal{A}_n^{MHV} = \mathcal{A}_n^{tree}W(\mathcal{C}_n) \quad (52)$$

with necessary identification of parameters. Since $\mathcal{A}_n^{tree}$ is rational function of kinematical variables it does not contribute to the cut. Hence we can rewrite optical theorem as

$$\begin{aligned} 2Im[W(\mathcal{C}_n)] &= 2Im\left(\frac{\mathcal{A}(in \rightarrow out)}{\mathcal{A}^{tree}(in \rightarrow out)}\right) \\ &= \sum_{states} \frac{\mathcal{A}^*(out \rightarrow all)\mathcal{A}(in \rightarrow all)}{\mathcal{A}^{tree}(in \rightarrow out)} \end{aligned} \quad (53)$$


At one loop level we have sum over products of tree-level amplitudes. Denoting this sum divided by $\mathcal{A}_n^{tree}$ as V_W (which can be found in appendix B) we have

$$Im[W(\mathcal{C})] = \int_{\mathcal{C}_L \mathcal{C}_R} V_W(\mathcal{C}_L, \mathcal{C}_R) \quad (54)$$



where integration goes over contours which one could get by breaking the loop into two parts, inserting special vertex, which one could find from summing over states in $\mathcal{N} = 4$ SYM and then by integration over contours which are limited by momentum conservation and light-like condition for every edge.

On the other hand the problem of finding imaginary part can be considered at diagrammatical level where the connection with the box in dual dimension makes it possible to apply Cutkosky rules. Of course, on this way there is no any summation over states. It would be nice to understand how the vertices from the dual amplitude picture occur in the game. The following picture arises if one considers the quadruple cut



It is interesting to note the role of the coefficient

$$\mathcal{C}_{MHV}^{2me} = \delta^8(\sum_{i=1}^n \lambda_i \eta_i) \Delta \quad (55)$$

which appears from quadruple cut of the MHV amplitude and is defined by the structure of tree amplitudes in $\mathcal{N} = 4$ SYM. In the Wilson loop calculation which is

blind to trees it appears while we go down from $D = 6$ to $D = 4$ dimensions, namely

$$z_0 \sim \frac{1}{\Delta} \quad (56)$$

7 On the geometry of UV/IR divergences

Let us discuss the interpretation of the divergent contributions. The IR singularity of the amplitude corresponds to the UV singularity of the cusps hence the very issue of the proper IR regularization of the amplitude is essentially related to the smoothing of the cusps in the polygon in the momentum space.

Let us make a few comments concerning the proper identification of the cusp anomaly in the geometrical terms [25, 26]. Since the amplitude is expressed in terms of the hyperbolic volumes and area in 3D AdS space it is natural to question what the cusp anomaly corresponds to in the same setting. That is we can start with the box with all external momenta off-shell which is finite. Then approaching on-shell limit for two external momenta the geometrical volume and area start to diverge which corresponds to the divergence of the Feynman diagram. Nevertheless we expect that the initial geometry is partially seen in the divergent terms.

Recall that $\Gamma_{cusp}(\theta, \alpha)$ is the cusp anomalous dimension which for the cusp angle θ at one loop behaves as

$$\Gamma_{cusp}(\theta, \alpha) = \frac{\alpha C_F}{\pi} (\theta \coth \theta - 1) + O(\alpha^2) \quad (57)$$

It turns out [25] that one-loop expression is nothing but the transition amplitude in AdS_3

$$\Gamma_{cusp}(\theta) \propto \langle v' | 1/\Delta_{S_3} | v \rangle \quad (58)$$

where two light-like vectors v and v' cross at the angle, and Δ_{S_3} is the corresponding Laplace operator on the $SU(2)$ group manifold. That is the one-loop anomaly can be attributed to the amplitude along the single edge of the basic simplex upon the analytic continuation [25]. Note that these geodesics connecting two vertices are dressed by the quadratic fluctuations.

Since the quantum geometry of the AdS_3 is governed by the $SL(2, C)$ Chern-Simons theory the corresponding Wilson loop is just the particle moving in this background. It is also possible to make the link with the AdS_2 geometry since the one-loop cusp anomaly can be written as the wave functional in the two dimensional YM theory on the disc integrated over its area.

$$\Gamma_{cusp}(\theta) \propto \int dA (Z(U, A) - Z(U, 0)) \quad (59)$$

where A is the area of the disc, U is the boundary holonomy and $Z(U, A)$ is partition function of the 2D YM theory in the disc geometry.

Since it is expected that the reparametrization of the boundary enters the answer it is natural to search for the Liouville interpretation of the cusp anomaly. Contrary to the finite contribution where the reparametrization part decouples and does not depend on the kinematical invariants we expect that divergent "Liouville" contribution has nontrivial kinematical dependence.

The possible arguments which however deserve more justification look as follows [26]. Consider two dimensional scalar field theory with the equation of motion

$$(\partial_t^2 - \partial_x^2)\phi + m^2\phi = 0 \quad (60)$$

whose solution has the following mode expansion

$$\phi(x, t) = \int \frac{d\beta}{2\pi} (a^*(\beta) e^{-im(x \sinh \beta - t \cosh \beta)} + a(\beta) e^{im(x \sinh \beta - t \cosh \beta)}) \quad (61)$$

It is convenient to introduce Rindler coordinates

$$\begin{aligned} x &= r \cosh \theta, & t &= r \sinh \theta \\ -\infty < \theta < +\infty & & 0 < r < +\infty \end{aligned} \quad (62)$$

in the space-time region $x > |t| > 0$. Upon the following Laplace transform with respect to the radial coordinate

$$\lambda_\theta(\alpha) = \int dr e^{imr \sinh \alpha} \left(-\frac{1}{r} \partial_\theta + im \cosh \alpha \right) \phi(r, \theta) \quad (63)$$

the commutation relation for the Laplace transformed field reads as

$$[\lambda(\alpha_1), \lambda(\alpha_2)] = i\hbar \tanh(\alpha_1 - \alpha_2)/2 \quad (64)$$

and the Hilbert space is spanned by vectors $a(\beta_n) \dots a(\beta_1)|vac >$ where vacuum state is defined as

$$a(\beta)|vac > = 0 \quad < vac|a^+(\beta) = 0 \quad (65)$$

One can introduce two-point function on the "rapidity plane"

$$F(\alpha_1 - \alpha_2) = < vac| \lambda(\alpha_1) \lambda(\alpha_2) |vac > \quad (66)$$

and explicit calculation amounts to the following answer [42]

$$F(\alpha - i\pi) = -\frac{1}{\pi} \alpha/2 \coth(\alpha/2) + \text{singular terms} \quad (67)$$

Hence the singular terms cancel in the difference $F(\alpha - i\pi) - F(-i\pi)$ which coincides with the cusp anomaly in agreement with the interpretation of [25] in the first quantized picture.

The relation with the Liouville model becomes clear upon the proper limiting procedure. To this aim we can try to represent Klein- Gordon equation of motion as the zero curvature condition for $SL(2, R)$ connection. Similarly the equation of motion in the Liouville model

$$(\partial_t^2 - \partial_x^2)\phi + \frac{m^2}{b}e^{b\phi} = 0 \quad (68)$$

can be considered as zero curvature condition for $SL(2, R)$ valued connection A_θ, A_r . It is convenient to introduce the monodromy matrix in the Liouville model

$$\mathbf{T}^\theta(\alpha) \propto e^{imR \sinh \alpha \sigma_3} \mathcal{P} \exp\left(\int dr A_r(r, \theta, \alpha)\right) \quad (69)$$

where R is cutoff, which defines $\lambda_{Liouv}(x)$ via relation

$$\lambda(\alpha) = -i \ln T_{11}(\alpha) \quad (70)$$

The latter reduces to the corresponding Klein-Gordon function involved into the cusp anomaly and in the weak coupling limit $b \rightarrow 0$ [42]

$$\lambda_{Liouv}(\alpha) \rightarrow \frac{b}{4} \lambda_{KG}(\alpha) \quad (71)$$

8 Discussion

In this paper we have discussed the different aspects of the duality between the calculation of the Wilson polygons and amplitudes in SUSY gauge theories focusing mainly on the one loop correspondence. It turns out that the duality for the MHV amplitude can be explicitly derived in the one-loop case. The derivation is remarkably simple and involves only the proper change of the variables and the relation between the Feynman integrals in the different space-time dimensions. The Wilson polygon to some extent can be thought as placed in the space of the Feynman parameters and it is in this space the change of variables is important. It was shown that the UV behavior of the Wilson polygon precisely maps into IR behavior of the amplitude which explains the correspondence between the regularizations observed earlier.

The change of variables found works well for the MHV amplitude only which can be expressed in terms of two mass-easy box diagrams and the generalization of the duality for the NMHV cases is required. Note that we have identified the key feature of the MHV kinematics - only in this case the integration over reparametrizations is decoupled which is not true for the rest of the cases. Therefore one could expect for the generic kinematics the emergence of the correlators of the Liouville modes responsible for the reparametrizations of the boundary contour. We consider the similar duality for the three-point function with one external particle on-shell. It was shown that

the duality can be formulated upon the insertion of the peculiar vertex operator into the Wilson triangle. We consider this example as providing the possible way for the generalization of the duality for the polarization-sensitive case. Let us emphasize that SUSY was not essentially used in our one-loop derivation of the duality. Probably the duality can be similarly developed for the non-SUSY theories as well.

It is worth to make more general comment concerning the relation of our analysis with the moduli space geometry. In the approach of [17, 18] the Schwinger parameters get mapped generically into the radial coordinate in AdS_5 and the moduli space of the complex structures $M_{g,n}$ where n is related to the number of the external legs in the amplitude. That is the Schwinger parametrization is closely related to the B model. On the other hand in our paper we exploited the picture with the emergent Kahler moduli which happens in A model. In principle one could imagine that a kind of the mirror transform on the level of the Feynman diagrams can be formulated and it would be very interesting to investigate this issue further. Note also that the A model under consideration allows the target space effective description in terms of the effective noncommutative gauge theory [41]. We hope to discuss the possible relation between the Wilson loops in $D = 6$ we have discussed with the corresponding object in the effective target space $D = 6$ gauge theory elsewhere.

The duality implies that a kind of the unitary technique can be developed for the calculation of the Wilson polygon as well. We have formulated the cut procedure for the one-loop Wilson polygon which involves the integration along the cut with the peculiar vertex-like operator. Along this line of reasoning we have also derived the analogue of the Landau equations for the singularities for the Wilson polygon in terms of the Feynman parameters. Let us emphasize that the geometry behind the Landau equations has a lot in common with the generic hyperbolic geometry behind the one-loop amplitudes. Actually the generic off-shell box diagram calculates the hyperbolic volume of the simplex defined by the kinematical invariants that is all divergences emerging upon some external particle tends to be on-shell are expected to carry some geometrical information about the initial hyperbolic geometry. We have shown that at the one-loop level this happens indeed.

In this paper we have discussed the one loop case only hence it would be very interesting to extend this analysis to the higher loops. The approach to the all-loop answer based on the quantum geometry of the momentum space suggested in [24] could be useful. Another promising development concerns the relation with the geometry of the knots which emerges because of the relation with the volumes of the hyperbolic spaces identified with the knot complements.


We are grateful to G. Korchemsky, N. Nekrasov, Yu. Makeenko and A. Rosly for the useful discussions. The work was supported in part by grants PICS- 07-0292165(A.G') and 09-02-00308(A.G., A.Z.). A.G. thanks FTPI at University of Minnesota where the part of the work was done for the kind hospitality and support. A.Z. thanks ICTP at Trieste where the part of the work was done for the kind

hospitality and support.

Appendix A Connection of scalar integrals in different dimensions

Here we briefly explain the connection between the scalar integrals in different dimensions [15]. Suppose, we have the following scalar integral

$$I^N(D; \nu_k) \equiv -i\pi^{-\frac{D}{2}}(\mu^2)^\epsilon \int d^D l \frac{1}{A_1^{\nu_1} A_2^{\nu_2} \dots A_N^{\nu_N}} \quad (72)$$



Than it can be shown that

$$I^N(D-2; \nu_k) = \sum_{i=1}^N z_i I^N(D-2; \nu_k - \delta_{ki}) + (D-1 - \sum_{j=1}^N \nu_j) z_0 I^N(D; \nu_k) \quad (73)$$

where

$$\sum_{i=1}^N (r_i - r_j)^2 z_i = 1 \quad (74)$$

$$z_0 = \sum_{i=1}^N z_i$$

In the main body of the text we choose $D = 6 + 2\epsilon$, $N = 4$, $\nu_i = 1$.

Appendix B Sum over states in terms of dual superconformal invariants

It is convenient to use $\mathcal{N} = 4$ on-shell formulation of $\mathcal{N} = 4$ SYM, in which all states are encoded in one super-wavefunction

$$\begin{aligned} \Phi(p, \eta) &= G^+(p) + \eta^A \Gamma_A(p) + \frac{1}{2} \eta^A \eta^B S_{AB}(p) + \frac{1}{3!} \eta^A \eta^B \eta^C \epsilon_{ABCD} \tilde{\Gamma}^D(p) \\ &+ \frac{1}{4!} \eta^A \eta^B \eta^C \eta^D \epsilon_{ABCD} G^-(p) \end{aligned} \quad (75)$$

We are interested in cuts of superamplitudes for n particles

$$\mathcal{A}_n(\lambda, \tilde{\lambda}, \eta) = \mathcal{A}_n(\Phi_1, \dots, \Phi_n) \quad (76)$$

Using of superamplitudes formalism one can easily obtain particular configuration of states using known projectors. As usually we use the two-component spinor formalism, where $p_i^{\alpha\dot{\alpha}} = \lambda_i^\alpha \tilde{\lambda}_i^{\dot{\alpha}}$ and $\langle i|j \rangle = \lambda_i^\alpha \lambda_{j\alpha}$.

While the $N^k\text{MHV}$ amplitude can be presented in terms of nested sums which are quite cumbersome expressions for MHV and NMHV cases are pretty simple

$$\mathcal{A}_n^{MHV} = \frac{\delta^{(4)}(\sum_{i=1}^n \lambda_i^\alpha \tilde{\lambda}_i^{\dot{\alpha}}) \delta^{(8)}(\sum_{i=1}^n \lambda_i^\alpha \eta_i^A)}{\langle 1|2 \rangle \dots \langle n-1|n \rangle \langle n|1 \rangle} \quad (77)$$

where the second (Grassmann) delta-function makes the supersymmetry manifest.

$$\mathcal{A}_n^{NMHV} = \frac{\delta^{(4)}(\sum_{i=1}^n \lambda_i^\alpha \tilde{\lambda}_i^{\dot{\alpha}}) \delta^{(8)}(\sum_{i=1}^n \lambda_i^\alpha \eta_i^A)}{\langle 1|2 \rangle \dots \langle n-1|n \rangle \langle n|1 \rangle} \sum_{(i,j)} R_{k;ij} \quad (78)$$

and all indexes are understood in the following way $i + n \equiv i$. Then $k + 2 \leq i < j \leq n + k - 1$ and $j - i \geq 2$. $R_{k;ij}$ are dual conformal invariants which are given by the following expressions

$$R_{k;ij} = \frac{\langle i|i-1 \rangle \langle j|j-1 \rangle \delta^{(4)}(\Xi_{k;ij})}{x_{ij}^2 \langle k|x_{ki}x_{ij}|j \rangle \langle k|x_{ki}x_{ij}|j-1 \rangle \langle k|x_{kj}x_{ji}|i \rangle \langle k|x_{kj}x_{ji}|i-1 \rangle} \quad (79)$$

Here the $\Xi_{k;ij}$ is

$$\Xi_{k;ij} = \langle k|x_{ki}x_{ij}|\theta_{jk} \rangle + \langle k|x_{kj}x_{ji}|\theta_{ik} \rangle \quad (80)$$

thus in this language R depends on $n - 2$ momenta.

For supermomentum delta-functions we will widely use the following identity [31] which could be easily proved

$$\delta^{(8)}(I)\delta^{(8)}(J) = \delta^{(8)}(I + J)\delta^{(8)}(J) \quad (81)$$

The summation over states is equivalent to the integration over $\int d^4\eta_{cut}$. For particular cuts, it was done in term of MHV vertex expansion in [30]. Here we present some results in simple forms using the language of dual conformal invariants R .

Thus, summation over states in L-loop cut is obtained by


$$\mathcal{A}_{cut}^{sum} = \int d^4\eta_{cut,1} d^4\eta_{cut,2} d^4\eta_{cut,L} d^4\eta_{cut,L+1} \mathcal{A}_{left}^{tree} * \mathcal{A}_{right}^{tree} \quad (82)$$

For $N^{L-1}\text{MHV} \times \text{MHV}$ cuts using (81) we can interpret the summation over states as the action of projection operators.

The result for $N^{L-1}\text{MHV} \times \text{MHV}$ cut, thus, reads as

$$\mathcal{A}_{cut}^{sum} = \frac{\delta^{(8)}(ext)}{\langle l_L | l_{L+1} \rangle^4} A_{N^{L-1}\text{MHV}split}^{gluons,tree}(++ \dots + ext; -- \dots - loop) A_{MHV,right}^{gluons,tree}(- - + \dots + loop; + \dots + ext) \quad (83)$$

If we cut MHV diagram at one loop (suppose loop momenta are l_1 and l_2) then for state sum we have



$$\mathcal{A}_{cut}^{sum} = \frac{\delta^{(8)}(ext)}{\langle l_1 | l_2 \rangle^4} A_{MHV}^{gluons,tree}(++ \dots + ext; -- loop) A_{MHV,right}^{gluons,tree}(- loop; + \dots + ext) \quad (84)$$

which agrees with formulas obtained in the literature.

To our purposes we need divide it on the tree-level amplitude which we have cut. That would be the operator which glues together Wilson loops:

$$\begin{aligned}
V_W &= \frac{\Pi_{ext} \langle i|i+1 \rangle}{\langle l_1|l_2 \rangle^4} A_{MHV}^{gluons,tree}(++\dots+_{ext}; --_{loop}) A_{MHV,right}^{gluons,tree}(-_{loop}; +\dots+_{ext}) \\
&= \frac{\langle i|i-1 \rangle \langle j|j-1 \rangle}{\langle l_1|j \rangle \langle l_1|j-1 \rangle \langle l_2|i \rangle \langle l_2|i-1 \rangle} < l_1|l_2 >^2
\end{aligned} \tag{85}$$

Another case of special interest for us is anti-MHV $\times$ MHV cut when we obtain

$$\mathcal{A}_{cut}^{sum} = \frac{\delta^{(8)}(ext)}{\langle l_L|l_{L+1} \rangle^4} A_{MHV}^{gluons,tree}(++_{ext}; --\dots-_{loop}) A_{MHV,right}^{gluons,tree}(-_{loop}; +\dots+_{ext}) \tag{86}$$

References

- [1] E. Witten, “Perturbative gauge theory as a string theory in twistor space,” Commun. Math. Phys. **252**, 189 (2004) [arXiv:hep-th/0312171].
- [2] L. F. Alday and J. M. Maldacena, “Gluon scattering amplitudes at strong coupling,” JHEP **0706** (2007) 064 [arXiv:0705.0303 [hep-th]].
- [3] A. M. Polyakov, “The wall of the cave,” Int. J. Mod. Phys. A **14**, 645 (1999) [arXiv:hep-th/9809057].
J. McGreevy and A. Sever, “Planar scattering amplitudes from Wilson loops,” JHEP **0808** (2008) 078 [arXiv:0806.0668 [hep-th]].
- [4] J. M. Drummond, G. P. Korchemsky and E. Sokatchev, “Conformal properties of four-gluon planar amplitudes and Wilson loops,” Nucl. Phys. B **795**, 385 (2008) [arXiv:0707.0243 [hep-th]].
J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “On planar gluon amplitudes/Wilson loops duality,” Nucl. Phys. B **795**, 52 (2008) [arXiv:0709.2368 [hep-th]].
- [5] A. Brandhuber, P. Heslop and G. Travaglini, “MHV Amplitudes in N=4 Super Yang-Mills and Wilson Loops,” Nucl. Phys. B **794**, 231 (2008) [arXiv:0707.1153 [hep-th]].
- [6] Z. Bern, L. J. Dixon, D. A. Kosower, R. Roiban, M. Spradlin, C. Vergu and A. Volovich, “The Two-Loop Six-Gluon MHV Amplitude in Maximally Supersymmetric Yang-Mills Theory,” arXiv:0803.1465 [hep-th].
- [7] J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “Hexagon Wilson loop = six-gluon MHV amplitude,” arXiv:0803.1466 [hep-th].
- [8] J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “Dual superconformal symmetry of scattering amplitudes in N=4 super-Yang-Mills theory,” arXiv:0807.1095 [hep-th].

- [9] J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “Conformal Ward identities for Wilson loops and a test of the duality with gluon amplitudes,” arXiv:0712.1223 [hep-th].
- [10] Z. Komargodski, “On collinear factorization of Wilson loops and MHV amplitudes in $N=4$ SYM,” JHEP **0805**, 019 (2008) [arXiv:0801.3274 [hep-th]].
L. F. Alday and J. Maldacena, “Minimal surfaces in AdS and the eight-gluon scattering amplitude at strong coupling,” arXiv:0903.4707 [hep-th].
- [11] L. F. Alday and R. Roiban, “Scattering Amplitudes, Wilson Loops and the String/Gauge Theory Correspondence,” arXiv:0807.1889 [hep-th].
- [12] N. Berkovits and J. Maldacena, “Fermionic T-Duality, Dual Superconformal Symmetry, and the Amplitude/Wilson Loop Connection,” arXiv:0807.3196 [hep-th].
N. Beisert, R. Ricci, A. Tseytlin and M. Wolf, “Dual Superconformal Symmetry from $AdS_5 \times S^5$ Superstring Integrability,” arXiv:0807.3228 [hep-th].
- [13] J. M. Drummond, J. M. Henn and J. Plefka, “Yangian symmetry of scattering amplitudes in $N=4$ super Yang-Mills theory,” arXiv:0902.2987 [hep-th].
- [14] O. V. Tarasov, “Reduction of Feynman graph amplitudes to a minimal set of basic integrals,” Acta Phys. Polon. B **29**, 2655 (1998) [arXiv:hep-ph/9812250].
- [15] G. Duplancic and B. Nizic, “Reduction method for dimensionally regulated one-loop N -point Feynman integrals,” Eur. Phys. J. C **35**, 105 (2004) [arXiv:hep-ph/0303184].
- [16] G. Duplancic and B. Nizic, “Dimensionally regulated one-loop box scalar integrals with massless internal lines,” Eur. Phys. J. C **20**, 357 (2001) [arXiv:hep-ph/0006249].
- [17] R. Gopakumar, “From free fields to AdS,” Phys. Rev. D **70**, 025009 (2004) [arXiv:hep-th/0308184].
- [18] O. Aharony, Z. Komargodski and S. S. Razamat, “On the worldsheet theories of strings dual to free large N gauge theories,” JHEP **0605**, 016 (2006) [arXiv:hep-th/0602226].
O. Aharony, J. R. David, R. Gopakumar, Z. Komargodski and S. S. Razamat, “Comments on worldsheet theories dual to free large N gauge theories,” Phys. Rev. D **75**, 106006 (2007) [arXiv:hep-th/0703141].
R. Gopakumar, “From free fields to AdS. III,” Phys. Rev. D **72**, 066008 (2005) [arXiv:hep-th/0504229].
- [19] A. Gorsky and V. Lysov, “From effective actions to the background geometry,” Nucl. Phys. B **718**, 293 (2005) [arXiv:hep-th/0411063].

- [20] A. I. Davydychev and R. Delbourgo, “A geometrical angle on Feynman integrals,” *J. Math. Phys.* **39**, 4299 (1998) [arXiv:hep-th/9709216].
- [21] A. I. Davydychev and J. B. Tausk, “A Magic connection between massive and massless diagrams,” *Phys. Rev. D* **53**, 7381 (1996) [arXiv:hep-ph/9504431].
- [22] N. I. Usyukina and A. I. Davydychev, “New results for two loop off-shell three point diagrams,” *Phys. Lett. B* **332**, 159 (1994) [arXiv:hep-ph/9402223].
- [23] Z. Bern, L. J. Dixon and V. A. Smirnov, “Iteration of planar amplitudes in maximally supersymmetric Yang-Mills theory at three loops and beyond,” *Phys. Rev. D* **72**, 085001 (2005) [arXiv:hep-th/0505205].
- [24] A. Gorsky, ”Quantum geometry of the momentum space and amplitudes in the N=4 SYM theory”, In Proceedings of ”Quarks-2008*” (May 2008) and memorial ”Landau-100” (June 2008) Conferences
A. Gorsky, To appear
- [25] A. V. Belitsky, A. S. Gorsky and G. P. Korchemsky, “Gauge / string duality for QCD conformal operators,” *Nucl. Phys. B* **667**, 3 (2003) [arXiv:hep-th/0304028].
- [26] A. Gorsky, “Spin chains and gauge / string duality,” *Theor. Math. Phys.* **142**, 153 (2005) [*Teor. Mat. Fiz.* **142**, 179 (2005)] [arXiv:hep-th/0308182].
- [27] D. J. Broadhurst, “Solving differential equations for 3-loop diagrams: Relation to hyperbolic geometry and knot theory,” arXiv:hep-th/9806174.
- [28] J.Drummond, J.Henn, G.Korchemsky, E.Sokatchev, [hep-th/0808.0491];
J. M. Drummond, J. Henn, G. P. Korchemsky and E. Sokatchev, “Generalized unitarity for N=4 super-amplitudes,” arXiv:0808.0491 [hep-th].
- [29] J. M. Drummond and J. M. Henn, “All tree-level amplitudes in N=4 SYM,” arXiv:0808.2475 [hep-th].
- [30] H. Elvang, D. Z. Freedman and M. Kiermaier, “Recursion Relations, Generating Functions, and Unitarity Sums in N=4 SYM Theory,” arXiv:0808.1720 [hep-th].
- [31] G. Georgiou, E. W. N. Glover and V. V. Khoze, “Non-MHV Tree Amplitudes in Gauge Theory,” *JHEP* **0407**, 048 (2004) [arXiv:hep-th/0407027].
- [32] A. M. Polyakov, “String theory and quark confinement,” *Nucl. Phys. Proc. Suppl.* **68**, 1 (1998) [arXiv:hep-th/9711002].
Y. Makeenko and P. Olesen, “Implementation of the Duality between Wilson loops and Scattering Amplitudes in QCD,” *Phys. Rev. Lett.* **102**, 071602 (2009) [arXiv:0810.4778 [hep-th]].

- [33] L. D. Landau, “On analytic properties of vertex parts in quantum field theory,” Nucl. Phys. **13**, 181 (1959).
- [34] R. E. Cutkosky, “Singularities and discontinuities of Feynman amplitudes,” J. Math. Phys. **1** (1960) 429.
- [35] R. Eden, P. Landshoff, D. Olive, J. Polkinghorne, “The analytic S-matrix,” (1966);
- [36] V. Smirnov, “Evaluating Feynman integrals,” Tracts Mod.Phys.211:1-244, (2004)
- [37] Z. Bern, L. J. Dixon and D. A. Kosower, “All next-to-maximally helicity-violating one-loop gluon amplitudes in $\mathcal{N} = 4$ super-Yang-Mills theory,” Phys. Rev. D **72**, 045014 (2005) arXiv:0412210 [hep-th].
- [38] A.Davydychev, A. I. Davydychev, “A Simple formula for reducing Feynman diagrams to scalar integrals,” Phys. Lett. B **263**, 107 (1991).
- [39] C. Anastasiou, A. Brandhuber, P. Heslop, V. Khoze, B. Spence, G. Travaglini, “Two-Loop Polygon Wilson Loops in N=4 SYM,” arXiv:0902.2245 [hep-th].
- [40] K. Hikami, “Generalized Volume Conjecture and the A-Polynomials: The Neumann-Zagier Potential Function as a Classical Limit of Quantum Invariant,” J. Geom. Phys. **57**, 1895 (2007) [arXiv:math/0604094].
T. Dimofte, S. Gukov, J. Lenells and D. Zagier, “Exact Results for Perturbative Chern-Simons Theory with Complex Gauge Group,” arXiv:0903.2472 [hep-th].
- [41] A. Iqbal, N. Nekrasov, A. Okounkov and C. Vafa, “Quantum foam and topological strings,” JHEP **0804**, 011 (2008) [arXiv:hep-th/0312022].
- [42] S. L. Lukyanov, “Correlators of the Jost functions in the Sine-Gordon model,” Phys. Lett. B **325**, 409 (1994) [arXiv:hep-th/9311189].